

AN ALGORITHM TO DETECT BLENDS WITH ECLIPSING BINARIES IN PLANET TRANSIT SEARCHES

HENK HOEKSTRA^{1,2,3}, YANQIN WU³ & ANDRZEJ UDALSKI⁴
Draft version February 2, 2008

ABSTRACT

We present an algorithm that can detect blends of bright stars with fainter, un-associated eclipsing binaries. Such systems contaminate searches for transiting planets, in particular in crowded fields where blends are common. Spectroscopic follow-up observations on large aperture telescopes have been used to reject these blends, but the results are not always conclusive. Our approach exploits the fact that a blend with a eclipsing binary changes its shape during eclipse.

We analyze original imaging data from the Optical Gravitational Lensing Experiment (OGLE), which were used to discover planet transit candidates. Adopting a technique developed in weak gravitational lensing to carefully correct for the point spread function which varies both with time and across the field, we demonstrate that ellipticities can be measured with great accuracy using an ensemble of images. Applied to OGLE-TR-3 and OGLE-TR-56, two of the planetary transit candidates, we show that both systems are blended with fainter stars, as are most other stars in the OGLE fields. Moreover, while we do not detect shape change when TR-56 undergoes transits, TR-3 exhibits a significant shape change during eclipses. We therefore conclude that TR-3 is indeed a blend with an eclipsing binary, as has been suggested from other lines of evidence. The probability that its shape change is caused by residual systematics is found to be less than 0.6%.

Our technique incurs no follow-up cost and requires little human interaction. As such it could become part of the data pipeline for any planetary transit search to minimize contamination by blends. We briefly discuss its relevance for the Kepler mission and for binary star detection.

Subject headings: binaries:eclipsing – stars: planetary systems

1. INTRODUCTION

Discovering planets outside the solar system is one of the key goals of modern astronomy. Since the first detection (Mayor & Queloz 1995) using the radial velocity technique, we have come to know of the existence of ~ 140 extra-solar planets. While radial velocity monitoring of nearby stars remains the most successful technique in this venture, a promising alternative is slowly gaining ground. This so-call ‘transit’ method focuses on detecting planets that transit their host stars. It requires continuous observing of a large number of stars, but can provide independent information concerning planet characteristics otherwise unobtainable by the radial velocity technique. A couple dozens planet transit searches are currently underway (see Horne 2003 for a review). Among these, the Optical Gravitational Lensing Experiment (OGLE) has announced a large number of planetary transit candidates (Udalski et al. 2002a; and follow-ups), and a number of these candidates have been confirmed spectroscopically (e.g., Konacki et al. 2003a, 2004a, 2004b; Bouchy et al. 2004; Pont et al. 2004; Torres et al. 2005). These are selected to resemble close-in gaseous planets (period ≤ 10 days) transiting main-sequence stars in the galactic disk. Many of the OGLE candidates (e.g. OGLE-TR-56, ~ 1.2 days) have orbital periods a factor of 2 shorter than the closest planets discovered by the radial velocity technique. The questions arise whether this is a new population of planets

and why they are not seen by the radial velocity method. If confirmed to be genuine planets, they pose intriguing challenges for understanding planet formation, migration and survival.

What types of objects can masquerade as planet transits? The success of OGLE in detecting planet transits relies partly on the extreme crowdedness and hence large base numbers in its fields. However, this advantage also brings on the masqueraders – a faint eclipsing binary system can project coincidentally (or in some cases, associate physically) near a brighter disk star. The deep eclipses and the ellipsoidal variations from the binary are then diluted by the light from the brighter star into shallow eclipses and little variations out of transit, mimicking the signatures of a transiting planet.

Sirko & Paczynski (2003) carefully studied the light-curves of these candidates and concluded that on average $\sim 50\%$ of these are contaminations by eclipsing binaries, with the shorter-period ones more likely to be so. Spectroscopic follow-up of a large number of these candidates (Konacki et al. 2003a; Dreizler et al. 2003) also reached a similar conclusion, though at a much greater observational expense. Moreover, spectroscopic observations are not always able to separate the blends from genuine planetary objects as the blended main star may show little or no velocity variations (see, e.g. Torres et al. 2004). High-quality photometric light-curves can be used to rule out

¹ Department of Physics and Astronomy, University of Victoria, Victoria, BC, V8P 5C2, Canada

² CITA, University of Toronto, Toronto, Ontario M5S 3H8, Canada

³ Department of Astronomy & Astrophysics, University of Toronto, 60 St. George Street, Toronto, Ontario M5S 3H8, Canada

⁴ Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland

the blends (Seager & Mallen-Ornelas 2003), but such data are difficult to obtain for the crowded OGLE fields. It is also possible to exclude some blending configurations by comparing the observed light-curves against synthetic light-curves constructed using model isochrones (Torres et al. 2004, 2005). This latter technique is more powerful if the blend and the main star are physical triples and therefore are likely coeval.

Our aim in this work is to provide an independent new method to recognize blends. Our method is efficient, assembly-line in style, and robust. It uses original imaging data and does not require any follow-up work. As such, this method may be broadly adopted in light of the fact that OGLE and other transiting searches are likely to produce an increasing number of planet candidates in the future. Moreover, our method is more suitable for detecting blends that are not physically associated (coincidental alignment) and thereby complements the light-curve method of Torres et al. (2004, 2005).

We propose to use the fact that a blended system, albeit unresolved in the images, always leaves a tell-tale sign: the shapes of their images are not round. The magnitude of the ellipticity depends on the angular separation and the relative brightness between the primary star and the secondary blend. As we show below, we can measure the shape of a typical blend in the OGLE fields with great precision. Comparison of the shape in and out of transit allows us to identify blends with eclipsing binaries. For instance, a star blended with an eclipsing binary with an undiluted eclipsing depth of 50% is expected to exhibit a factor of 2 change in its ellipticity between the two phases. The actual change in shape may be smaller, though still detectable, as a typical star in the OGLE fields is multiply blended.

The success of this technique depends critically on how well we measure the shape of a star, in relation to other stars in the same image. This is where the only major obstacle in this method arises: the point-spread-function (PSF) varies across the image due to a multitude of distortions in the photon pathway. It also varies with time as the pathway changes and the seeing fluctuates. PSF anisotropy and seeing change the shapes of the objects and renders raw measurements of the ellipticity unreliable. A similar problem exists in weak gravitational lensing, where one has to disentangle the lensing induced distortions in the shapes of faint galaxies from these observational effects. Fortunately, the weak lensing community has studied this problem in great detail and has come up with solutions which we adapt to the case in hand. We note that the method we develop here has aspects unique to the stellar problem.

Among the hundreds of transiting systems published by the OGLE-III team (Udalski et al. 2002a, 2002b, 2002c), we choose to focus our initial efforts on two candidates, OGLE-TR-3 and OGLE-TR-56. On the basis of spectroscopic follow-up observations with 8m class telescopes, these two candidates were identified as likely planetary candidates since their host stars show little or no velocity variations (Konacki et al. 2003a; Dreizler et al. 2003). TR-56 undergoes genuine flat-bottom transit and has detectable radial velocity variations, both consistent with a planet explanation. For this candidate, the blending sce-

nario was examined in detail by Torres et al. (2005) who were able to confirm the planetary nature of this object using a combined analysis of the light curve and radial velocity measurements. The interpretation for TR-3, however, is more open to debate. It shows no significant velocity variations, its light curve contains hints of a secondary eclipse as well as out-of-eclipse fluctuations (Sirko & Paczynski 2003; Konacki et al. 2003a). The method presented here provides a completely independent assessment of the identities of these two objects.

We briefly describe the data in §2. The shape measurement technique is described in detail in §3. In §4 we provide an extensive test of our analysis and present the results for the two planet transit candidates in §5.

2. DATA

The data we analyze were obtained during the third phase of OGLE (OGLE III, Udalski et al. 2002a). These were collected using the 1.3m Warsaw telescope at the Las Campanas Observatory, equipped with the 8k MOSAIC camera. The field of view of the camera is about 35 by 35 arcminutes, with a pixel scale of $0''.26/\text{pixel}$. The observations were done in the *I*-band, and have exposure times of 120s.

Our analysis does not require the full field, so instead we use small cuts of 600 by 600 pixels, not necessarily centered on the target candidate. For TR-3 we have 109 images in-transit and 308 images out-of-transit, whereas we have 65 and 259 images, respectively, for TR-56. We retrieved all in-transit images, which results in a broad range in seeing. To minimize the systematic errors caused by the seeing correction (see §3.2), we have selected out-of-transit images such that their seeing distribution resembles that of the in-transit data. This is illustrated in Figure 1.




FIG. 1.— Histogram of the seeing distribution of the images used in the analysis for TR-3 (left) and TR-56 (right). The solid lines indicate the distribution for the in-transit frame, whereas the dotted histogram corresponds to the out-of-transit data. The seeing distributions of the out-of-transit data were matched to resemble the in-transit distribution.

3. METHOD

In this section we discuss the shape measurements, focussing on how to deal with the variable PSF. The methodology is based on the techniques developed for weak gravitational lensing applications (e.g., see Kaiser, Squires & Broadhurst 1995; Hoekstra et al. 1998), and we adopt their notation. The correction for the PSF can be split into two separate steps. The first one is the correction for

the anisotropic part of the PSF, which induces an ellipticity in addition to the intrinsic ellipticity of the object under investigation. The second step is the correction for the circularization by the PSF (i.e., seeing), which typically lowers the ellipticity. For both steps, we require a set of comparison stars which can be presumed to be intrinsically round.

To quantify the shapes, we use the central second moments I_{ij} of the image fluxes and form the two-component polarisation

$$e_1 = \frac{I_{11} - I_{22}}{I_{11} + I_{22}} \text{ and } e_2 = \frac{2I_{12}}{I_{11} + I_{22}}. \quad (1)$$

Because of photon noise, unweighted second moments cannot be used. Instead we use a circular Gaussian weight function, with a dispersion r_s :

$$I_{ij} = \int d^2\vec{x} f(\vec{x}) x_i x_j W(|x|, r_s). \quad (2)$$

where x_i is the pixel number in the direction of the i -axis, pointing away from the centroid of the object. For the weight function W we adopt a Gaussian with a dispersion r_s . For the analysis presented here, the weighted moments are measured from the images within an aperture with a radius of 6 pixels, and we take $r_s = 1.5$ pixels, which is the optimal width for a seeing of $0''.9$. These choices suppress the contributions from nearby stars.

3.1. Anisotropic PSF

In practice, the PSF will not be isotropic. Instead, the images are typically convolved with an anisotropic PSF, which induces coherent ellipticities in the images. In order to recover the true “shape” of the blend, we need to undo the effect of the PSF anisotropy. The correction scheme we use is based on that developed by Kaiser, Squires & Broadhurst (1995), with modifications described in Hoekstra et al. (1998).

The effect of an anisotropic PSF on the polarisation e_α of an object is quantified by the “smear polarisability” P_{ii}^{sm} , which measures the response of the polarisation to a convolution with an anisotropic PSF, and can be estimated for each object from the data (see Hoekstra et al. 1998 for the correct expressions).

Having measured the polarisations and smear polarisabilities, the corrected polarisations are given by

$$e_i^{\text{cor}} = e_i^{\text{obs}} - P_{ii}^{\text{sm}} p_i, \quad (3)$$

where p_i is a measure of the PSF anisotropy. It is measured using a true point source by

$$p_i = e_i^{\text{PSF}} / P_{ii}^{\text{sm,PSF}}, \quad (4)$$

where e_i^{PSF} are the measured ellipticity of the point source and $P_{ii}^{\text{sm,PSF}}$ the diagonal components of its smear polarisability tensor. Formally, the correction requires the use of the full two by two tensor, but the off-diagonal terms are typically small. Examination of the measured values indicates that they are consistent with noise. We therefore only use the diagonal terms in the correction for PSF anisotropy.

This correction has been tested extensively in the case of galaxies convolved with an anisotropic PSF (e.g., Hoekstra et al. 1998; Erben et al. 2001). For this application, the correction works well, because galaxies are centrally concentrated, and their shapes are well characterized by the quadrupole moments.

In the case of two or more nearby point sources the situation is somewhat different: the shape is not well described by a simple quadrupole, and higher order moments are expected to contribute to the polarisation. To explore this in more detail, we created images that were convolved with a Moffat (1969) profile and then convolved with a line (which simulates the PSF anisotropy). A detailed discussion of this study can be found in the Appendix. Here we summarize the main conclusion.

The simulations indicate that the correction given by Eqn. 3 is incomplete and that an additional term proportional to $|p|$ (the total size of the anisotropy) is needed. This leads to an improved correction for PSF anisotropy, albeit empirical, given by

$$e_i^{\text{cor}} = e_i^{\text{obs}} - P_{ii}^{\text{sm}} p_i - \alpha \sqrt{p_1^2 + p_2^2}, \quad (5)$$

where the value of α depends on the configuration of the point sources (separation and flux ratio) and the seeing. We found that the size of α is proportional to the polarisation of the object. This is supported by an examination of the residuals in the shapes of the objects in the OGLE data.

The fact that α is proportional to the polarisation is not surprising: when the polarisation is larger, the higher order moments become more important. However, in the case of OGLE, blends with more than one source are likely. Consequently, it is difficult to compute the expected value of α . Instead, we determine the value empirically by fitting a term proportional to $|p|$ to the shape measurements.

The PSF anisotropy depends on the position of the object on the chip and it typically varies with time. Fortunately, it is possible to characterize the spatial variation of the PSF anisotropy with a low order polynomial model fitted to a subsample of the objects identified as suitable stars (i.e., the stars should be bright but not saturated). This works particularly well for the data used here, as we use relatively small regions around the OGLE transit candidates. For the analysis here we model the spatial variation by a second order polynomial. Such a model is derived for each exposure and used to undo the effect of the PSF anisotropy.

The derivation of the PSF anisotropy model implicitly assumes that the set of comparison stars are intrinsically round: i.e., the observed polarisation is solely caused by PSF anisotropy. It is possible to reject wide separation binaries (or blends) from this set on the basis of their large ellipticities, but it is more difficult to reject stars that have a small intrinsic ellipticity because of a companion. However, so long as the number of comparison stars is sufficiently large, because their position angles are uncorrelated with each other and with the PSF anisotropy, we still can obtain an unbiased model for the PSF anisotropy.




FIG. 2.— Residuals in p_1 (upper panels) and p_2 (lower panels) for the comparison stars are plotted here (points with error bars) against position on the chip. They are obtained from the measured values (points without error bars) after subtracting a second-order polynomial for the PSF anisotropy. This simple correction leaves no detectable trend in p_i . The scatter may be partially related to the intrinsic ellipticity arising from blending.

In the case of OGLE, severe crowding means most bright stars are blended with fainter stars. As a result, the noise introduced by the blends can be substantial. Nevertheless, we can find a set of brighter stars which are comparatively less affected by blending and provide good estimates for the PSF anisotropy. The result of this procedure carried over one frame is presented in Figure 2: we detected significant PSF anisotropy, and found that a second-order polynomial is sufficient to remove the PSF anisotropy across the whole image, leaving residual p_1 and p_2 scattering randomly around the zero-level.

In the absence of blending and shot noise, the corrected ellipticities should all be zero (assuming the model used to correct for the PSF anisotropy is perfect). However, blending gives rise to non-zero ellipticities for the stars and is partly responsible for the residual anisotropy in Figure 2. We measure this “intrinsic” ellipticity of the stars used in the PSF anisotropy correction using repeated observations taken by the OGLE team (as this procedure reduces the shot noise). We then subtract the “intrinsic” ellipticities from the observed ones and obtain an improved fit. We found that this iteration had little effect on the results, because of the random orientations of the blends.

Large values of PSF anisotropy are typically a nuisance, as they imply larger corrections. However, the large range of PSF anisotropy (Fig. 3) exhibited by the OGLE observations is helpful for the purpose of our paper: it allows us to examine the accuracy of the correction for PSF anisotropy in more detail, and to understand the limitation of our algorithm.




FIG. 3.— Histograms for the measured PSF anisotropy, p_1 and p_2 (averaged over all selected stars in an image), for TR-3 (upper panels) and TR-56 (lower panels) frames. The solid lines are for the in-transit frames, whereas the dotted lines correspond to the out-of-transit measurements. The distributions for the in and out-of-transit frames span similar ranges.

3.2. Correction for seeing variation

The second step in our correction procedure is to account for the effect of seeing, i.e., the isotropic part of the PSF. Typically, an object will appear rounder with increasing seeing. An example is presented in the left panels of Figure 4, which shows the ellipticities for one of the stars in TR-3 field as the seeing varies. The dependence on seeing can be rather complicated, with some configurations appearing more eccentric with increasing seeing.

In the simple case of a single blend, one could use simulations to attempt to determine the seeing dependence. This is not feasible here, because the target stars are on average blended with 1.5 objects. Instead, we make use of the fact that the observations span a large range in seeing (see Fig. 1 and Fig. 4) to remove the seeing dependence empirically. Using the following model, individually tuned for each star,

$$e_\alpha = a_0 + a_1 \cdot \text{seeing} + a_2 \cdot \text{seeing}^2. \quad (6)$$

This second-order fitting is sufficient to remove any visible seeing dependence for, e.g., the object shown in Figure 4. From now on, we report the shape measurement for a fiducial seeing, taken to be 1 arcsecond.

4. TESTING THE ALGORITHM

We selected a sample of a total of 171 stars around the transit candidate in the two fields. These stars have some range in brightness and ellipticity. We demonstrate below our capabilities in removing the effects of PSF anisotropy and seeing.




FIG. 4.— *Left panels*: variation of the two components of the polarisation with seeing for an elongated object in the TR-3 field, after correcting for PSF anisotropy. The polarisation decreases with increasing seeing, as most stars do. *Right panels*: the value of the polarisation at a fiducial seeing of 1 arcsecond after we account for the variation with seeing using a second-order polynomial model. Note the different vertical scales for e_1 and e_2 . The error bars are determined from the scatter in the polarisations after correcting for seeing. Although determined independently, the errors for e_1 and e_2 are comparable.

4.1. Correcting for PSF anisotropy

To examine the accuracy of the PSF anisotropy correction, we split the in-transit data into two subsets of similar sizes: one with large PSF anisotropy ($|p|$) and one with small $|p|$. The two subsets have a similar range in seeing.

In Figure 5, we present the differences in ellipticity measurements for these comparison stars, without correcting for PSF anisotropy (upper panels), after using equation (3) to correct for the anisotropy (middle panels), and after using equation (5) to correct for the anisotropy (lower panels).⁵ This experiment convinced us that we can remove PSF anisotropy successfully from our data. The reduced χ^2 for the results in the lower panels of Figure 5 is close to unity, indicating that the estimated errors are a fair estimate of the statistical uncertainty in the measurements.

In producing Figure 5, we have applied a seeing correction that is based on the combined in-transit data, minimizing the systematics caused by the latter correction. As mentioned above, the seeing ranges are similar for both samples, so PSF anisotropy is the only systematic relevant for comparison. Upper panels in Figure 6 expand the view from the lower panels of Figure 5 for the small ellipticity objects, while lower panels in Figure 6 shows results from the same procedure using out-of-transit data. For some of the brighter objects, the achieved error bars are as small as $\sim 1.2 \times 10^{-4}$. This capability to measure shapes accurately brings about another potential application for the algorithm described here: finding binary stars that are too close, or are too different in fluxes, to be resolved (also see

⁵ This additional correction is tiny for objects with small ellipticities.

§6).




FIG. 5.— *Upper panels*: the difference in e_1 (left) and e_2 (right) between the in-transit data with large PSF anisotropy and the in-transit data with small anisotropy, before correction for PSF anisotropy. The offsets from zero simply reflects the effects of the PSF anisotropy. *Middle panels*: the difference between the two samples when Eqn. 3 is used to correct for PSF anisotropy. A clear trend $\propto e_\alpha$ can be discerned. Note that the vertical scale has been decreased by a factor two. *Lower panels*: the results when the empirical correction given by Eqn. 5 is applied for the PSF anisotropy correction. No trend with the shape of the object is visible. The vertical scale has been decreased by another factor of two. The error bars have been determined from a bootstrap resampling of the data.




FIG. 6.— *Upper panels*: differences in e_1 (left) and e_2 (right) between the sample with large and small PSF anisotropy for the in-transit observations, limited to the region of small ellipticity. *Lower panels*: the same differences, but now between two samples taken from the out-of-transit observations.

4.2. Correcting for Seeing

We now examine the reliability of the correction for seeing. If this is successful, we will be able to accurately measure the in and out-of-transit ellipticity changes in planet candidates, and constrain the blending scenario.

As shown in Figure 1, the seeing distributions for the in and out-of-transit data were chosen to be similar, i.e. we selected out-of-transit images such that the two distributions match. This approach minimizes the sensitivity of our results to systematic errors caused by the adopted seeing correction. We fit equation (6) to the in and out-of-transit measurements separately. The resulting differences between the in and out-of-transit data are presented in Figure 7. Panels a and b show the differences in e_1 and e_2 respectively.

The lower panels in Figure 7 show histograms of the differences in units of the estimated measurement uncertainty. For a normal distribution, this should be a Gaussian with a dispersion of 2, which is indicated by the solid smooth line. This Gaussian provides a fair match to the observed scatter, but the data show more outliers than what would be expected from a normal distribution. For Δe_1 we obtain a reduced $\chi^2_{\text{red}}=1.31$ and for Δe_2 we find a similar value of $\chi^2_{\text{red}}=1.37$, larger than the expected value around unity. These values reduce to 1.14 and 1.16 respectively when we reject objects that are more than 3σ away from zero.

The bootstrap analysis provides an estimate of the random error but not of the systematic error. The results presented in Figure 7 suggest that the estimated errors are typically correct, but in a few cases, residual systematic errors are still present in the data leading to the excess of outliers. Nevertheless, the results presented in Figure 7 do suggest that for most objects we can measure the difference in shapes between the in and out-of-transit data accurately. We have attempted to identify what is causing some of the outliers, but have not been able to find an obvious way to improve the measurements. We suspect that it might be due to imperfections in the correction for PSF anisotropy. We also note that some of the objects are not present in all exposures (as they lie too close to the edge), which might lead to differences in the actual seeing distributions, which in turn can lead to systematic errors in the shape differences.

The size of the final error bar as used in Figure 7 depends on the number of frames used as well as on the apparent magnitude of the object: the shape measurements in a single frame will be noisier for fainter stars. This is demonstrated in Figure 8, which shows the error in e_1 and e_2 as a function of apparent magnitude for the out-of-transit shape measurements.⁶ As expected, the errors increase with magnitude. This is more clearly seen for “rounder” objects, which are affected less by the correction for seeing and the last term in equation 5 for the PSF anisotropy correction.

We also computed the smallest possible error bar as a function of apparent magnitude using simulated images. In these images, which have the same noise properties as the OGLE data, we measured the scatter in the shape of a point source. In this case, the error is solely due to Poisson noise. The result is given by the dashed curve in Figure 8.

The actual error bars are larger, because of the uncertainties introduced by the empirical corrections for PSF anisotropy and seeing. Finally, Figure 8 also demonstrates that the accuracy with which one can measure shapes is excellent: the typical uncertainty for a star with $m_I = 14$ is $\sim 3 \times 10^{-4}$.




FIG. 7. — *panel a*: difference Δe_1 between the out-of-transit and in-transit measurement of the polarisation for various stars in the TR-3 and TR-56 fields (note that the transit candidates are not included in this histogram). *panel b*: similar to panel a, but for Δe_2 . For both components the results suggest that the shape difference can be measured with great accuracy. *panel c*: histogram of the number of objects as a function of the ratio between Δe_1 and the corresponding measurement error. *panel d*: the same, but for Δe_2 . If the errors follow a normal distribution, this should resemble a Gaussian with a dispersion of 2, which is indicated by the smooth curve. The agreement with a normal distribution is fair, but a larger than expected number of outliers is found, suggestive of residual systematics. As before, the error bars were obtained from a bootstrap resampling of the data.

5. APPLICATION TO TRANSIT CANDIDATES

The results presented in the previous section demonstrates our ability to accurately measure the shapes of objects in the OGLE fields. In this section we present results for the two OGLE planet transit candidates. Table 1a lists the final polarisations for TR-3 and TR-56 at a fiducial seeing of 1 arcsecond measured from the out-of-transit images. We detect a significant polarisation for both transit candidates, thus implying that they are both blended with other sources. In fact, most stars studied in the crowded OGLE fields show evidence of blending (or even multiple blending). Within a circle of $1''$ radius, an average star is surrounded by 0.6 companions, with a mean flux ratio of 4% and a mean separation of $0.7''$.

The resulting average ellipticity of the blend depends mainly on the brightness of the primary star: the brighter the star, the smaller the ellipticity. Figure 9 shows the distribution of e_1 and e_2 for the analysed stars in the fields of TR-3 and TR-56 (indicated by the crosses). The two tran-

⁶ Although the errors in both components of the polarisation are derived separately, they typically are comparable.

sit candidates are indicated by the open circles, with TR-3 being the point on the left. Although the distribution is peaked towards round objects, the observed ellipticities for the transit candidates are by no means anomalous.




FIG. 8.— Derived errors in e_1 and e_2 as a function of apparent magnitude of the object. The solid points correspond to objects with $|e| < 0.015$, i.e., “rounder” objects, whereas the open circles indicate the errors for objects with $|e| > 0.015$. The error bars are larger for more elongated objects because of the larger uncertainty introduced by the empirical corrections for PSF anisotropy and seeing. The dashed line indicates the theoretical minimum value of the error bar as a function of magnitude. This curve is computed from simulated images of a point source, with noise properties corresponding the OGLE observations.

To test whether the ‘planet-like’ transit is caused by a blended eclipsing binary, we also list in Table 1b differences in shape between the out and in transit data for these two stars: a significant change in shape would confirm that the blend is an eclipsing binary.

TABLE 1

Results for TR-3 and TR-56, corrected for PSF anisotropy for a fiducial seeing of 1 arcsecond.

(a)	e_1	e_2
TR-3	-0.0234 ± 0.0005	-0.0052 ± 0.0005
TR-56	0.0101 ± 0.0005	-0.0005 ± 0.0005
(b)	$e_1(out) - e_1(in)$	$e_2(out) - e_2(in)$
TR-3	-0.0020 ± 0.0005	-0.0007 ± 0.0005
TR-56	-0.0001 ± 0.0006	0.0000 ± 0.0006

We detect no change in shape in TR-56, suggesting that the observed transit is a genuine planetary transit, in line with evidences from radial velocity, line-curve analysis and isochrone fitting (e.g., Torres et al. 2005).

In TR-3, however, we do observe a change in shape: the ellipticity in-transit is larger. The errors inferred from the bootstrap analysis suggest a significance of 4.2σ . However, the results presented in Section 4.2 and Figure 7 indicate

that the distribution of errors is not exactly Gaussian, but has tails. We therefore need to account for the possibility that the change in shape is caused by residual systematics. To this end, a more conservative estimate of the significance of the change in shape for TR3 can be obtained by considering the fraction of studied objects that show a difference at least as large as TR-3.




FIG. 9.— The crosses indicate the values of e_1 and e_2 for the additional stars that have been analysed in the out-of-transit images of TR-3 and TR-56. The measurement errors on the shapes are negligible and have not been displayed. The open circles indicate the results for TR-3 (left) and TR-56 (right). The distribution of points is peaked towards round objects, but few objects have truly no detectable ellipticity. The observed ellipticities for the transit candidates are by no means anomalous.

Of the objects in the fields of TR-3 and TR-56, accurate shapes could be determined for 171 of them. None of these objects show an ellipticity change as large as TR-3, and we can only derive a lower limit to the probability for the observed shape change in TR-3 to be caused by systematic effects: the probability is less than $1/171 \sim 0.6\%$. This is larger than the probability of a $4-\sigma$ event (0.03%) but still sufficiently small for us to conclude that it is very likely that TR-3 is indeed a blend with an eclipsing binary system.

If TR-3 is only singly blended, its polarisation should decrease by a factor of 2 during eclipse, an effect that should be eminently detectable. However, we find that its ellipticity increases by $\sim 10\%$ during eclipse. This can be explained if TR-3 is multiply blended. In fact, we have also measured TR-3’s polarization (out-of-transit) using different weight functions (r_s in eq. [2]). It varies with r_s differently than a singly-blended object would, suggesting that it is indeed multiply blended. In the case of multiple blending, provided that the primary star is much brighter than the blending stars, the resulting ellipticity is given by

$$e_i^{\text{obs}} \approx \sum_{j=1}^N e_{i,j} \quad (7)$$

where N is the number of blends, and $e_{i,j}$ is the contri-

bution from each blend. Consequently, if two blends have opposite signs for e_i , the polarisation can actually increase during an eclipse.

If we assume that the observed eclipse is caused by a blend with an eclipsing binary, with a eclipse depth 50% (i.e., a full eclipse of an equal mass system), we can place limits on the configuration of the blend. To do so, we note that the depth of the observed transit is 2%, which implies that the flux of the presumed binary contributes 4% to the total flux. Under these assumptions, the change in ellipticity indicates that the binary is located 0.38 ± 0.06 arcseconds from the brighter star. If the eclipse depth is reduced to 25%, the presumed binary contributes 8% of the flux instead, and the separation decreases to 0.26 ± 0.04 arcseconds. These numbers are below the resolution limit of the photometry (by analysing centroid shift in and out of eclipse, $\sim 1''$), and the blend is likely a background source (as opposed to a physical triple with the main star).

Interestingly, by examining the light-curve in detail, Konacki et al. (2003b) have also come to a similar conclusion that TR-3 is likely a blend of a background eclipsing binary with a foreground bright star. Our result here confirms their suggestion and predicts the position of the blend. The separation from the main star is small but should be detectable by HST observations.

6. CONCLUSIONS

We have presented an algorithm that can detect blends of bright stars with fainter eclipsing binaries. Such systems contaminate searches for transiting planets, in particular in crowded fields where blends are common. This technique provides a cheap way to find such blends, thus minimizing the amount of time required on large aperture telescope for spectroscopic follow-up of planet candidates.

We have demonstrated the accuracy with which shapes can be measured using imaging data from the Optical Gravitational Lensing Experiment (OGLE). Our method requires a careful correction of the point spread function which varies both with time and across the field. To this end we have adopted a method developed in weak gravitational lensing with modifications necessary for this particular application.

We have tested the correction for PSF anisotropy in great detail, using a sample of 171 stars surrounding the two planet transit candidates studied here. Comparison of samples with large and small PSF anisotropy indicates that this correction can be applied with great accuracy. For a star with an apparent magnitude $m_I = 14$, we obtain a 1σ uncertainty of $\sim 3 \times 10^{-4}$ in the polarisation.

Applied to OGLE-TR-3 and OGLE-TR-56, two of the planetary candidates, we show that both systems are indeed blended with fainter stars, as are most other stars in the OGLE fields. In the case of TR-56 we do not detect a change in shape in and out of transit, consistent with it indeed being a genuine planetary object. For TR-3 we observe a significant change in shape. If we adopt the error bars from the bootstrap analysis, the significance is 4.2σ . However, the distribution of errors is not precisely Gaussian, but has tails. A more conservative estimate of the significance, estimated from the observed distribution of

shape differences, provides an upper limit of 0.006 to the probability that the observed change is caused by residual systematics. Our results favour the scenario where TR-3 is caused by a blend with a background eclipsing binary, in line with evidences from other studies.

A number of studies have appeared since the OGLE announcement of transit candidates, mostly aiming at distinguishing blends from genuine planets. In contrast to some of these studies which carry out follow-up spectroscopy using large telescopes, our approach uses original imaging data and is a value-added application. Moreover, unlike studies which perform detailed light-curve fitting or isochrone stellar model fitting, our method is assembly-line in style and can be applied to a large number of transit candidates without too much human interaction. Lastly, our technique is especially suited to finding blends that are not physically associated with the bright star,⁷ and is therefore complementary to the isochrone fitting technique which is more powerful for the physical triple case.

Given the efficiency in dealing with a large number of objects without requiring additional data, the shape method may also be useful for other planetary transit searches, in particular the NASA Kepler mission. This transit mission aims to detect $\sim 10^3$ giant inner planets and $\sim 10^2$ terrestrial planets. Recently the target survey area has been moved to a higher galactic latitude to reduce the confusion by blends with eclipsing binaries. A quick examination of the USNO-B catalogue (Monet et al. 2003) in this new field suggests that the stellar density is ~ 8 times less dense than that in the OGLE field, with a similar number distribution in stellar magnitudes. However, stars in Kepler have a PSF of $6''$ radius, we therefore expect each bright star ($m_v < 14$) to have ~ 2.4 companions within the PSF envelope, compared to 0.6 ($< 1''$) in the OGLE case. The probability of blending with an eclipsing binary is likely enhanced by a similar ratio. More study is necessary to determine the false-positive rate due to blending in Kepler, armed with the experience from OGLE. Nevertheless, we expect that our shape technique can be readily applied to this mission.

The achieved accuracy in measuring the shape of stars also bodes well for another potential application of our algorithm: finding binary stars that are too close to be resolved, yet too far apart for radial velocity studies. By detecting small deviations from circularity, we should be able to discover intermediate separation binaries ($\sim 10 - 1000 AU$) with flux ratio as low as 1%, within a large volume of our galaxy. This will not only complement existing binary searches, but its high efficiency may also disclose binary population with an unprecedented rate such as to enable new and meaningful statistical studies. In a subsequent paper we will investigate this application in more detail, and apply it to wide field imaging data from the EXPLORE project (Mallén-Ornelas et al. 20003; Yee et al. 2003), which were obtained with the aim of finding transiting planets.

AU acknowledges support from the Polish KBN grant 2P03D02124 and the grant ‘‘Subsydium Profesorskie’’ of the Foundation for Polish Science.

⁷ These blends cause larger ellipticity.




FIG. 10.— (a) Measured polarisation as a function of PSF anisotropy p_α , for two point sources of equal flux, separated by $0''.4$ and a seeing of 1 arcsecond. The thin solid line indicates the values without any correction for PSF anisotropy. A single point source would show a linear trend, but because of the second point source the polarisation is constant for $p_\alpha > 0$. The dashed line corresponds to the polarisation when equation 3 is used to correct for PSF anisotropy. A clear trend $\propto |p|$ can be seen (in this case, the PSF anisotropy is given by p_α , with the other component zero). If we use equation 5, with an appropriate coefficient for the slope, we obtain the dotted line, which is effectively independent of PSF anisotropy. (b) The same as (a), but for a second component with a flux ratio of 0.5. The coefficient to obtain the dotted line is different from the one used in panel (a).

APPENDIX

IMPROVING THE CORRECTION FOR PSF ANISOTROPY

As indicated by Figure 5 the correction for PSF anisotropy using equation 3 leaves a systematic residual, roughly proportional to the polarisation. This correction scheme has been used extensively in weak lensing applications, and has been tested in great detail. The difference between the analysis of galaxies and the blends considered here, is that the shapes of galaxies are well characterized by their quadrupole moments. In the case of two point sources, higher order moments contribute to the moments.

In this section we examine how to improve the correction for PSF anisotropy, in particular we justify the use of equation 5. Unlike the case for galaxies (Kaiser et al. 1995), this problem is too complicated to solve analytically. Instead we study the effect of PSF anisotropy on simulated images of two point sources. We also note that new methods have been developed in which the images of the objects are decomposed into a series of localized basis functions. For instance, Bernstein & Jarvis (2002) use Laguerre expansions, whereas Refregier (2003) adopted weighted Hermite polynomials. The advantage of these methods might be that they can quantify higher order moments of the images. Nevertheless, as we will show below (and in §4), the empirical extension of the Kaiser et al. (1995) method is adequate for the results presented here.

We create well oversampled images of two point sources, and convolve these with a Moffat function, with a width given by the required seeing. These images are then convolved with a “line”, which simulates the effect of PSF anisotropy. Examples for two configurations are indicated by the thin solid lines in Figure 10. The results presented in this Figure are for a case where the PSF anisotropy is given by p_α alone, with the other component set to zero. A single point source would show a linear trend with p_α , but because of the second point source the slope changes when p_α changes sign.

The next step is to correct these polarisations for PSF anisotropy. If we use equation 3, we obtain the dashed lines in Figure 10. In both cases we see a clear residual $\propto |p_\alpha|$. More general simulations, with both components of the PSF anisotropy non-zero, indicate that the slope is actually $\propto |p|$, and not just $|p_\alpha|$. These results have led us to consider an additional term in the correction $\propto \sqrt{p_1^2 + p_2^2}$, leading to equation 5.

Based on a large set of simulations, we found that the slope of the trend is proportional to e_α . This is not completely surprising, because the amplitude of the polarisation is a measure of the importance of the second point source, and consequently a measure of the relevance of higher order moments. Hence the additional term in the correction for PSF anisotropy is given by $\gamma e_\alpha |p|$.

We examined what value for γ yields the best correction. The results indicate that γ depends on the configuration, in particular on the flux ratio. For instance, in Figure 10a we used $\gamma = 0.7$ and in Fig. 10b we obtained the best result for $\gamma = 0.6$ to obtain the improved corrections, indicated by the dotted lines. Overall, the range in γ appears to be fairly small, although its value is difficult to determine if the polarisations are small (i.e., when the additional correction is small).

We note that the conclusions listed above are based on our empirical study of the simulated images. using equation 5, the correction for PSF anisotropy works well for most of the situations (as is the case for the ones presented in Fig. 10). In some extreme cases, however, with large values for the polarisation and PSF anisotropy, the correction leaves significant residuals (relative errors as large as 10%).

REFERENCES

- Bernstein, G.M. & Jarvis, M. 2002, *AJ*, 123, 583
 Bouchy, F., Pont, F., Santos, N.C., Melo, C., Mayor, M., Queloz, D., & Udry, S. 2004a, *A&A*, 421, L13.
 Dreizler, S.; Hauschildt, P. H.; Kley, W.; Rauch, T.; Schuh, S. L.; Werner, K.; Wolff, B. 2003, *A&A*, 402, 791
 Erben, T., van Waerbeke, L., Bertin, E., Mellier, Y., & Schneider, P. 2001, *A&A*, 377 717
 Hoekstra, H., Franx, M., Kuijken, K. & Squires, G. 1998, *ApJ*, 504, 636
 Horne, K. 2003, *astro-ph/0301249*
 Kaiser, N., Squires, G., & Broadhurst, T. 1995, *ApJ*, 449, 26
 Konacki, M.; Torres, G.; Jha, S.; Sasselov, D. D. 2003a, *Nature*, 421, 507
 Konacki, M., Torres, G., Sasselov, D. D., & Jha, S. 2003b, *ApJ*, 597, 1076
 Konacki, M., Torres, G., Sasselov, D.D., Pietrzyński, G., Udalski, A., Jha, S., Ruiz, M.T., Gieren, W., & Minniti, D. 2004a, *ApJL*, 609, L37.
 Konacki, M., Torres, G., Sasselov, D.D., and Jha, S., 2004b, *ApJL*, in press, *astro-ph/0412400*.
 Mallén-Ornelas, G, Seager, S., Yee, H.K.C., Minniti, D., Gladders, M.D., Mallén-Fullerton, G.M., Brown, T.M. 2003, *ApJ*, 582, 1123
 Mayor, M.; Queloz, D. 1995, *Nature*, 378, 355
 Moffat, A.F.J. 1969, *A & A*, 3, 455
 Monet, D. G.; Levine, S. E.; Canzian, B. et al. 2003, *AJ*, 125, 984
 Pont, F., Bouchy, F., Queloz, D., Santos, N.C., Melo, C., Mayor, M., and Udry, S. 2004, *A&A*, 426, L15.
 Refregier, A. 2003, *MNRAS*, 338, 35
 Seager, S.; Mallén-Ornelas, G. 2003, *ApJ*, 585, 1038
 Sirko, E.; Paczynski, B. 2003, *ApJ*, 592, 1217
 Torres, G.; Konacki, M.; Sasselov, D. D.; Jha, S. 2004, *ApJ*, 614, 979
 Torres, G.; Konacki (2), M.; Sasselov, D. D.; Jha, S. 2005, *ApJ*, *astro-ph/0410157*
 Udalski, A., Paczynski, B., Zebruski, K., Szymanski, M., Kubiak, M., Soszynski, I., Szweczyk, O., Wyrzykowski, L., & Pietrzynski, G. 2002a, *Acta Astron.*, 52, 1
 ——— 2002b, *Acta Astron.*, 52, 115
 Udalski, A., Szweczyk, O., Zebrun, K., Pietrzynski, G., Szymanski, M., Kubiak, M., Soszynski, I., Wyrzykowski, L. 2002c, *Acta Astron.*, 52, 317
 Yee, H.K.C., Mallén-Ornelas, G, Seager, S., Gladders, M.D., Brown, T., Minniti, D., Ellison, S., & Mallén-Fullerton, G. 2003, to appear in the Proceedings of the SPIE conference: Astronomical Telescopes and Instrumentation, *astro-ph/0208355*